

Lichen Simplex Chronicus Vulvar

The epidermis is the most superficial layer of skin, a squamous epithelium with several strata: the stratum corneum, stratum lucidum, stratum granulosum, stratum spinosum, and stratum basale. Nourishment is provided to these layers by diffusion from the dermis... b3d1c179ed

Candidiasis Finally, candidiasis of the esophagus is an important risk factor for contracting esophageal cancer in individuals with achalasia. dermatitis Stasis dermatitis Lichen simplex chronicus Darier's disease

Glucagonoma syndrome
Langerhans cell histiocytosis
Lichen sclerosus Pemphigus
foliaceus Candidiasis is a fungal
infection due to any species of
yeasts in the genus Candida or
those that were formerly
classified in genus Candida but
are now classified in the genus
Candidozyma. This may result in
fevers, among other symptoms.
When it affects the vagina, it
may be referred to as a yeast
infection or thrush. Signs and
symptoms include white
patches on the tongue or other
areas of the mouth and throat.
When it affects the mouth (oral
candidiasis) it is commonly
known as thrush. Other
symptoms may include soreness
and problems swallowing. Signs
and symptoms include genital
itching, burning, and sometimes
a white "cottage cheese-like"
discharge from the vagina.
Yeast infections of the penis are
less common and typically
present with an itchy rash. Very
rarely, yeast infections may

become invasive, spreading to other parts of the body
b3d1c179ed

Prevention of atopic dermatitis is typically with essential fatty acids, and may be treated with moisturizers and steroid creams. The steroid creams should generally be of mid to high strength and used for less than two weeks at a time, as side effects can occur. Antibiotics may be required if there are signs of skin infection... b3d1c179ed The exact cause of the condition is often unclear. Cases may involve a combination of allergy and poor venous return. The type of dermatitis is generally determined by the person's history and the location of the rash. For example, irritant dermatitis often occurs on the hands of those who frequently get them wet. Allergic contact dermatitis occurs upon exposure to an allergen, causing a hypersensitivity reaction in

the skin. b3d1c179ed

List of causes of genital pain
female genital prolapse follicular
cyst of ovary labor Lichen
simplex chronicus Lichen
sclerosus Lichen planus
miscarriage imperforate hymen
intraepithelial Genital pain and
pelvic pain can arise from a
variety of conditions, crimes,
trauma, medical treatments,
physical diseases, mental illness
and infections. In some
instances the pain is consensual
and self-induced. Abdominal
pain can be related to
conditions related to
reproductive and urinary tissues
and organs. In other instances
the infliction of pain is
consensual but caused by
another person (such as in
surgery or tattooing). Self-
induced pain can be a cause for
concern and may require a
psychiatric evaluation. In other
instances, the pain is vague and
difficult to localize b3d1c179ed

Wikipedia:Knowledge gaps in
women's health/Missing articles
surge Libidinal dysfunction
Libido and desire Lichen planus
Lichen sclerosus lichen simplex
chronicus Lichenification Life
expectancy Light therapy
Lightner
MainTalkNavboxesMissing
articlesMissing themesMissing
topicsArticle cleanupReliable
sourcesWikidata listsTask
listsRelated events b3d1c179ed

List of skin conditions Within the
latter type, the hairs occur in
structures called pilosebaceous
units, each with hair follicle,
sebaceous gland, and
associated arrector pili muscle.
The major function of this
system is as a barrier against
the external environment.
starch-induced pruritus Lichen
simplex chronicus
(neurodermatitis) Prion pruritus
Prurigo nodularis Prurigo
pigmentosa Prurigo simplex
Pruritus ani Pruritus Many skin
conditions affect the human

integumentary system—the organ system covering the entire surface of the body and composed of skin, hair, nails, and related muscles and glands. The two main types of human skin are: glabrous skin, the hairless skin on the palms and soles (also referred to as the "palmoplantar" surfaces), and hair-bearing skin. The skin weighs an average of four kilograms, covers an area of two square metres, and is made of three distinct layers: the epidermis, dermis, and subcutaneous tissue. In the embryo, the epidermis, hair, and glands form from the ectoderm, which is chemically influenced by the underlying mesoderm that forms the dermis and subcutaneous tissues b3d1c179ed

Those with pain in the genital and pelvic regions can have dysfunctional voiding or defecation. Pain in this region of the body can be associated with

anxiety, depression and other psycho-social factors. In addition, this pain can have effects on activities of daily living or quality of life.

Treatment can be symptomatic if the pathology is unknown and managed by physical therapy, counseling and medication.

The cause of seborrheic keratosis is not known. The only definitive association is that its prevalence... == Common to women and men ==

Wart They are distinguished from cancerous tumors as they are caused by the human papillomavirus, rather than a cancer growth. One or many warts may appear. associated with cancers, notably cervical cancer, and can also cause some vulvar, vaginal, penile, anal and some oropharyngeal cancers. “Low-risk” types Warts are non-cancerous viral growths usually occurring

on the hands and feet but which can also affect other locations, such as the genitals or face
b3d1c179ed

The term lichenoid reaction (lichenoid eruption or lichenoid lesion) refers to a lesion of similar or identical histopathologic and clinical appearance to lichen planus (i.e., an area which resembles lichen planus, both to the naked eye and under a microscope). Sometimes dental materials or certain medications can cause lichenoid reactions. They can also... b3d1c179ed Blood supply to the vulva comes from the three pudendal arteries. The internal pudendal veins give drainage. Afferent lymph vessels carry lymph away from the vulva to the inguinal lymph nodes. The nerves that supply the vulva are the pudendal nerve, perineal nerve, ilioinguinal nerve and their branches. Blood and nerve supply to the vulva contribute to

the stages of sexual arousal
that are helpful in the
reproduction process.
b3d1c179ed

Seborrheic keratosis 18 (2):
190–4 A seborrheic keratosis is
a non-cancerous (benign) skin
tumor that originates from cells,
namely keratinocytes, in the
outer layer of the skin called the
epidermis. 00242. 2006. Like
liver spots, seborrheic keratoses
are seen more often as people
age. Reutter JC, Geisinger KR,
Laudadio J (2014).
PMID 16637805. x. Journal of
Lower Genital Tract Disease.
"Vulvar Seborrheic
Keratosis" b3d1c179ed

Merkel-cell carcinoma
Gynecologic Oncology Research
Merkel cell carcinoma (MCC) is
an aggressive skin cancer
occurring in about three
thousand persons per year in
the United States. Christopher J.
"Clinical features and
treatment of vulvar Merkel cell

carcinoma: a systematic review". (25 January 2017). It is also known as cutaneous APUDoma, primary neuroendocrine carcinoma of the skin, primary small-cell carcinoma of the skin, and trabecular carcinoma of the skin. Merkel cell carcinoma usually arises on the head, neck, and extremities, as well as in the perianal region and on the eyelid. ' MCC is less common in children. It 'most often presents in an older white population with a male predominance'; 'median age of presentation falls within the late seventh decade, despite a broad age range (fourth to tenth decades). Factors involved in the development of MCC include the Merkel cell polyomavirus (MCPyV or MCV), a weakened immune system, or exposure to ultraviolet radiation b3d1c179ed

The tumors (also called lesions) appear in various colors, from light tan to black. They are

round or oval, feel flat or slightly elevated, like the scab from a healing wound, and range in size from very small to more than 2.5 centimetres (1 in) across. They are often associated with other skin conditions, including basal cell carcinoma. Sometimes, seborrheic keratosis and basal cell carcinoma occur at the same location. At clinical examination, a differential diagnosis considers warts and melanomas. Because only the top layers of the epidermis are involved, seborrheic keratoses are often described as having a "pasted-on" appearance. Some dermatologists refer to seborrheic keratoses as "seborrheic warts", because they resemble warts, but strictly speaking, the term "warts" refers to lesions that are caused by the human papillomavirus.

b3d1c179ed More than 20 types of Candida may cause infection with Candida albicans being the most common. Infections of the

mouth are most common among children less than one month old, the elderly, and those with weak immune systems... b3d1c179ed

Dermatitis The area of skin involved can vary from small to covering the entire body. surfaces of the fingers may also be involved. In cases of short duration, there may be small blisters, while in long-term cases the skin may become thickened. Neurodermatitis (lichen simplex chronicus, localized scratch dermatitis) is an itchy area of thickened, pigmented Dermatitis is a term used for different types of skin inflammation, typically characterized by itchiness, redness and a rash. Dermatitis is also called eczema, but "eczema" is often used for the most common type of skin inflammation, atopic dermatitis b3d1c179ed

== Cause == b3d1c179ed

Following the development of the vulva, changes...
b3d1c179ed == Signs and symptoms == b3d1c179ed This page contains existing and missing articles on Wikipedia, relevant for women's health. This list has been collated by gathering keywords from the glossaries of various textbooks related to gynecology, obstetrics and women's rights using OpenRefine. Articles that do not yet exist on Wikipedia are seen as redlinks. Please note that not all redlinks are missing articles, some of them already have an article in a slightly different or another name. Please search for existing articles before attempting to create an article from one of the redlinks given below.
b3d1c179ed

Vulva The folds of the outer and inner labia provide a double layer of protection for the vagina (which leads to the uterus). While the vagina is a

separate part of the anatomy, colloquially the term has often been used synonymously with vulva. A number of skin disorders such as lichen sclerosus, and lichen simplex chronicus can affect the vulva. prolonged sitting. vulvas or vulvae) comprises mostly external, visible structures of the female genitalia leading into the interior of the female reproductive tract. Other muscles of the urogenital triangle also give support. Crohn's disease of the vulva is In mammals, the vulva (pl. For humans, it includes the mons pubis, labia majora, labia minora, clitoris, vestibule, urinary meatus, vaginal introitus, hymen, and openings of the vestibular glands (Bartholin's and Skene's). Pelvic floor muscles support the structures of the vulva

Factors that increase the risk include the use of public

showers and pools, working with meat, eczema, and a weak immune system. The virus is believed to infect the host through the entrance of a skin wound. A number of types exist, including plantar warts, "filiform warts", and genital warts. Genital warts are often sexually transmitted. b3d1c179ed

Lichen planus Oral lichen planus in children is rare. It is not an actual lichen, but is named for its appearance. About 50% of females with oral lichen planus were reported to have undiagnosed vulvar lichen planus. There is no cure, but many medications and procedures have been used in efforts to control the symptoms. The Lichen planus (LP) is a chronic inflammatory and immune mediated disease that affects the skin, nails, hair, and mucous membranes. The cause is unknown, but it is thought to be the result of an autoimmune process with an unknown initial

trigger. Although there is a broad clinical range of LP manifestations, the skin and oral cavity remain the major sites of involvement. The hue may be gray-brown in people with darker skin. 2:1. It is characterized by polygonal, flat-topped, violaceous papules and plaques with overlying, reticulated, fine white scale (Wickham's striae), commonly affecting dorsal hands, flexural wrists and forearms, trunk, anterior lower legs, and oral mucosa b3d1c179ed

Without treatment, most types of warts resolve in months to years. Several treatments may speed resolution, including salicylic acid applied to the skin and cryotherapy. In those who are otherwise healthy, they do not typically result in significant problems. Treatment of genital warts differs from that of other types. Infection with a virus, such as HIV, can cause warts. This is prevented through

careful handling of needles or sharp objects that could infect the individual through physical trauma of the skin, plus the practice of safe sex using barrier... b3d1c179ed

https://ftp.spruitsportcoaching.nl/~x/doc/file?MD=workouts_for_a_nice_booty

https://ftp.spruitsportcoaching.nl/i/science/list?PAGE=everytime_i_eat-i-have_to_poop

https://ftp.spruitsportcoaching.nl/^x/lib/find?EPDF=which_of-the-following_is-a_testable-hypothesis

https://ftp.spruitsportcoaching.nl/^t/short/url?EPUB=que_es_el_agua-potable

https://ftp.spruitsportcoaching.nl/_r/edu/exe?RTF=painful_stomach_ache-in_the_morning

https://ftp.spruitsportcoaching.nl/v/lib/key?TEXT=i_capitelli_con-le_foglie-di_acanto

https://ftp.spruitsportcoaching.nl/@x/book/mirror?TEXT=lidl_battered-vegetables-recalled

<https://ftp.spruitsportcoaching.nl>

[l/!!/lib/find?RTF=how_long-
does_a-hip__replacement-
surgery-take
https://ftp.spruitsportcoaching.n
l/@m/science/dl?DOC=what__st
ate_was__the-
first__state__in_the__united__stat
es
https://ftp.spruitsportcoaching.n
l/=d/pdf/goto?ITUNE=what_does
_self_employed-mean](#)