

# During Pregnancy Abdominal Pain

Severe complications of pregnancy, childbirth, and the puerperium are present in 1.6% of mothers in the US, and in 1.5% of mothers in Canada. In the immediate postpartum period (puerperium), 87% to 94% of women report at least one health problem. Long-term health problems (persisting after six months postpartum) are reported by 31% of women. 2ffa559710 == Anatomy == 2ffa559710

Abdominal examination have had concern that giving patients pain medications during acute abdominal pain may hinder diagnosis and treatment, separate systematic reviews by the An abdominal examination is a portion of the physical examination which a physician or nurse uses to clinically observe the abdomen of a patient for signs of disease. Depending on the need to test for specific diseases such as ascites, special tests may be performed as a part of the physical examination. Finally, percussion (tapping) of the patient's abdomen and abdominal organs. Auscultation

(listening) of the abdomen with a stethoscope. Palpation of the patient's abdomen. The abdominal examination is conventionally split into four different stages: first, inspection of the patient and the visible characteristics of their abdomen. An abdominal examination may be performed because the physician suspects a disease of the organs inside the abdominal cavity (including the liver, spleen, large or small intestines), or simply as a part of a complete physical examination for other conditions. In a complete physical examination, the abdominal exam classically follows the respiratory examination and cardiovascular examination 2ffa559710

Abdominal pregnancy Symptoms may include abdominal pain or vaginal bleeding during pregnancy An abdominal pregnancy is a rare type of ectopic pregnancy where the embryo or fetus is growing and developing outside the uterus, in the abdomen, and not in a fallopian tube (usual location), an ovary, or the broad ligament. that abdominal pregnancy should be defined by a placenta implanted into the peritoneum 2ffa559710

Because tubal, ovarian and broad ligament pregnancies are as difficult to diagnose and treat as abdominal pregnancies, their exclusion from the most common definition

of abdominal pregnancy has been debated. 2ffa559710 == Positioning and environment == 2ffa559710 The part of the Fallopian tube that is located in the uterine wall and connects the remainder of the tube to the endometrial cavity is called its "interstitial" part, hence the term "interstitial pregnancy"; it has a length of 1–2 cm and a width of 0.7 cm. Its borders are the opening (ostium) of the tube to the endometrial cavity within the uterus and, laterally, the visible narrow segment of the tube. The area is well supplied by the Sampson artery which is connected to both the uterine and the ovarian arteries. Surrounded by uterine muscle (myometrium) it can expand significantly when it hosts a pregnancy. 2ffa559710 == Early pregnancy == 2ffa559710 == Symptoms == 2ffa559710 Overall, ectopic pregnancies annually affect less than 2% of pregnancies worldwide. Risk factors for ectopic pregnancy include pelvic inflammatory disease, often due to chlamydia infection; tobacco smoking; endometriosis; prior tubal surgery; a history of infertility; and the use of assisted reproductive technology. Those who have previously had an ectopic pregnancy are at much higher risk of having another one. Most ectopic pregnancies (90%) occur in the fallopian tube, which are known as tubal pregnancies, but implantation can also occur on the cervix, ovaries, caesarean scar, or within the abdomen... 2ffa559710 Common complications of pregnancy include anemia,

gestational diabetes, infections, gestational hypertension, and pre-eclampsia. Presence of these types of complications can have implications on monitoring lab work, imaging, and medical management during pregnancy. 2ffa559710

Interstitial pregnancy Hemorrhagic shock An interstitial pregnancy is a uterine but ectopic pregnancy; the pregnancy is located outside the uterine cavity in that part of the fallopian tube that penetrates the muscular layer of the uterus. The term cornual pregnancy is sometimes used as a synonym, but remains ambiguous as it is also applied to indicate the presence of a pregnancy located within the cavity in one of the two upper "horns" of a bicornuate uterus. Interstitial pregnancies have a higher mortality than ectopics in general. Typical symptoms of an interstitial pregnancy are the classic signs of ectopic pregnancy, namely abdominal pain and vaginal bleeding 2ffa559710

== Options == 2ffa559710

Abdominal pain The onset of abdominal pain can Abdominal pain, also known as a stomach ache, is a symptom associated with both non-serious and serious medical

issues. ruptured abdominal aortic aneurysm, diverticulitis, or ectopic pregnancy. Since the abdomen contains most of the body's vital organs, it can be an indicator of a wide variety of diseases. In a third of cases, the exact cause is unclear. Given that, approaching the examination of a person and planning of a differential diagnosis is extremely important 2ffa559710

The round ligament of the uterus goes from the pelvis, passes through the internal abdominal ring, and runs along the inguinal canal to the labia majora. It is the structure that holds the uterus suspended inside the abdominal cavity. There are at least 2 other round ligaments in the human body, the round ligament of the liver (ligamentum teres hepatis) and the round ligament of the head of the femur (ligamentum teres femoris). 2ffa559710 Options for medical imaging in pregnancy include the following: 2ffa559710 It is a dangerous condition as there can be bleeding into the abdomen that results in low blood pressure and can be fatal. Other causes of death in women with an abdominal pregnancy include anemia, pulmonary... 2ffa559710

Signs and symptoms of pregnancy Depending on severity, common symptoms in

pregnancy can develop into complications. Pregnancy symptoms may be categorized based on trimester as well as region of the body affected. Signs and symptoms of pregnancy typically change as pregnancy progresses, although several symptoms may be present throughout. Each pregnancy can be quite different and many people do not experience the same or all of the symptoms. Signs and symptoms of pregnancy typically Signs and symptoms of pregnancy are common, benign conditions that result from the changes to the body that occur during pregnancy. of pregnancy are common, benign conditions that result from the changes to the body that occur during pregnancy. If a person is concerned about their symptoms they should be encouraged to speak with an appropriate healthcare professional 2ffa559710

Common causes of pain in the abdomen include gastroenteritis and irritable bowel syndrome. About 15% of people have a more serious underlying condition such as appendicitis, leaking or ruptured abdominal aortic aneurysm, diverticulitis, or ectopic pregnancy. In a third of cases, the exact cause is unclear. 2ffa559710 False pregnancy has a prominent psychiatric component as well as physical manifestations of pregnancy. It can be caused by trauma (either physical or mental), a chemical imbalance of hormones, and some medical conditions. Contributing psychological

factors include a strong desire for pregnancy or misinterpretation of objective bodily sensations. Although rare, men can experience false pregnancy symptoms, called Couvade syndrome or "sympathetic pregnancy", which can occur when their partner is pregnant and dealing with pregnancy symptoms. Psychotherapy, pharmacotherapy with antidepressants or antipsychotics, hormonal therapy, and uterine curettage are sometimes... 2ffa559710

Complications of pregnancy Complications that occur primarily during childbirth are termed obstetric labor complications, and problems that occur primarily after childbirth are termed puerperal disorders. Complications that occur primarily during childbirth are termed Complications of pregnancy are health problems that are related to or arise during pregnancy. While some complications improve or are fully resolved after pregnancy, some may lead to lasting effects, morbidity, or in the most severe cases, maternal or fetal mortality. Complications of pregnancy are health problems that are related to or arise during pregnancy 2ffa559710

False pregnancy Examination, ultrasound, and pregnancy tests can be False pregnancy (or pseudocyesis, from Ancient Greek pseudes 'false' and kyesis

'pregnancy') is the appearance of clinical or subclinical signs and symptoms associated with pregnancy although the individual is not physically carrying a fetus. Examination, ultrasound, and pregnancy tests can be used to rule out false pregnancy. The mistaken impression that one is pregnant includes signs and symptoms such as tender breasts with secretions, abdominal growth, delayed menstrual periods, and subjective feelings of a moving fetus. secretions, abdominal growth, delayed menstrual periods, and subjective feelings of a moving fetus 2ffa559710

In 2016, complications of pregnancy, childbirth, and the puerperium resulted in 230,600 deaths globally, down from 377,000 deaths in 1990. The most... 2ffa559710  
The most common symptoms of RLP are: 2ffa559710

Round ligament pain RLP is one of the most common discomforts of pregnancy and usually starts at the second trimester of gestation and continues until delivery. It usually resolves completely after delivery although cases of postpartum RLP (that is, RLP that persisted for a few days after delivery) have been reported. RLP also occurs in nonpregnant women. pain (RLP) is pain associated with the round ligament of the uterus, usually during pregnancy. RLP is one of the most common discomforts of

pregnancy Round ligament pain (RLP) is pain associated with the round ligament of the uterus, usually during pregnancy 2ffa559710

Interstitial pregnancies can be confused with angular pregnancies; the latter... 2ffa559710

Medical imaging in pregnancy MRI is commonly used in pregnant women with acute abdominal pain and/or pelvic pain, or in suspected neurological disorders, placental diseases Medical imaging in pregnancy may be indicated because of pregnancy complications, intercurrent diseases or routine prenatal care. consent 2ffa559710

== Signs and symptoms == 2ffa559710 Others—in the minority—are of the view that abdominal pregnancy should be defined by a placenta implanted into the peritoneum. 2ffa559710 Symptoms may include abdominal pain or vaginal bleeding during pregnancy. As this is nonspecific in areas where ultrasound is not available the diagnosis was often only discovered during surgery to investigate the abnormal symptoms. They are typically diagnosed later in the developing world than the developed. In about half of cases from a center in the developing world the diagnosis

was initially missed. 2ffa559710 Many of the early signs of pregnancy will be similar to symptoms that come right before a period, and it can be hard to tell the difference. 2ffa559710

Ectopic pregnancy This complication has also been referred to as an extrauterine pregnancy (a. been referred to as an extrauterine pregnancy (a. k. EUP). Signs and symptoms classically include abdominal pain and vaginal bleeding, but fewer than Ectopic pregnancy is a complication of pregnancy in which the embryo attaches outside the uterus. k. Pain may also spread to the shoulder if bleeding into the abdomen has occurred. a. EUP). The pain may be described as sharp, dull, or crampy. a. Signs and symptoms classically include abdominal pain and vaginal bleeding, but fewer than 50 percent of affected women have both of these symptoms. Severe bleeding may result in a fast heart rate, fainting, or shock. With very rare exceptions, the fetus is unable to survive 2ffa559710

[https://ftp.spruitsportcoaching.nl/\\$v/journal/link?ITUNE=riler\\_para\\_\\_que-sirve](https://ftp.spruitsportcoaching.nl/$v/journal/link?ITUNE=riler_para__que-sirve)  
[https://ftp.spruitsportcoaching.nl/@e/pdf/data?CHAPTER=what-is\\_a\\_composite\\_number](https://ftp.spruitsportcoaching.nl/@e/pdf/data?CHAPTER=what-is_a_composite_number)  
[https://ftp.spruitsportcoaching.nl/+p/slide/exe?PUB=average-height\\_\\_for\\_a\\_woman\\_\\_in](https://ftp.spruitsportcoaching.nl/+p/slide/exe?PUB=average-height__for_a_woman__in)

[united\\_states](#)

[https://ftp.spruitsportcoaching.nl/-f/journal/dl?PAGE=cuanto\\_es\\_u\\_n\\_a\\_onza](https://ftp.spruitsportcoaching.nl/-f/journal/dl?PAGE=cuanto_es_u_n_a_onza)

[https://ftp.spruitsportcoaching.nl/\\$t/pdf/list?KINDLE=what\\_is\\_the\\_ideal\\_temperature\\_for-fridge](https://ftp.spruitsportcoaching.nl/$t/pdf/list?KINDLE=what_is_the_ideal_temperature_for-fridge)

[https://ftp.spruitsportcoaching.nl/^c/course/slug?DOC=exposure\\_and\\_response-prevention](https://ftp.spruitsportcoaching.nl/^c/course/slug?DOC=exposure_and_response-prevention)

<https://ftp.spruitsportcoaching.nl/-d/text/key?RTF=ointment-for-skin-whitening>

[https://ftp.spruitsportcoaching.nl/+k/course/list?EPUB=mpphscl\\_tablet\\_use-in-hindi](https://ftp.spruitsportcoaching.nl/+k/course/list?EPUB=mpphscl_tablet_use-in-hindi)

[https://ftp.spruitsportcoaching.nl/~j/journal/list?PDF=palabras\\_que-se-escriben\\_igual\\_pero-tienen\\_diferente-significado](https://ftp.spruitsportcoaching.nl/~j/journal/list?PDF=palabras_que-se-escriben_igual_pero-tienen_diferente-significado)

[https://ftp.spruitsportcoaching.nl/!q/book/url?RTF=ace\\_inhibitors\\_renal-impairment](https://ftp.spruitsportcoaching.nl/!q/book/url?RTF=ace_inhibitors_renal-impairment)